

# Presynaptic Facilitation by Neuropeptide Signaling Mediates Odor-Driven Food Search

Cory M. Root,<sup>1</sup> Kang I. Ko,<sup>1</sup> Amir Jafari,<sup>1</sup> and Jing W. Wang<sup>1,\*</sup>

<sup>1</sup>Neurobiology Section, Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093, USA

\*Correspondence: [jw800@ucsd.edu](mailto:jw800@ucsd.edu)

DOI 10.1016/j.cell.2011.02.008

## SUMMARY

Internal physiological states influence behavioral decisions. We have investigated the underlying cellular and molecular mechanisms at the first olfactory synapse for starvation modulation of food-search behavior in *Drosophila*. We found that a local signal by short neuropeptide F (sNPF) and a global metabolic cue by insulin are integrated at specific odorant receptor neurons (ORNs) to modulate olfactory sensitivity. Results from two-photon calcium imaging show that starvation increases presynaptic activity via intraglomerular sNPF signaling. Expression of sNPF and its receptor (sNPFR1) in Or42b neurons is necessary for starvation-induced food-search behavior. Presynaptic facilitation in Or42b neurons is sufficient to mimic starvation-like behavior in fed flies. Furthermore, starvation elevates the transcription level of *sNPFR1* but not that of *sNPF*, and insulin signaling suppresses *sNPFR1* expression. Thus, starvation increases expression of *sNPFR1* to change the odor map, resulting in more robust food-search behavior.

## INTRODUCTION

The modulation of behavior by basic physiological need is essential for animal survival. Physiological modulation is often accomplished by release of neuromodulators that alter neuronal excitability or network properties (Destexhe and Marder, 2004). In particular, appetite and satiety modulate feeding behavior in most animals through the actions of neuropeptides. In mammals, the hypothalamus, an important brain region controlling appetite (Berthoud, 2002), integrates hormonal signals such as ghrelin, insulin, and leptin from the gut, pancreas, and adipose tissues, respectively. Activation of neurons containing neuropeptide Y (NPY) and AgRP in the arcuate nucleus of the hypothalamus augments food intake (for review, see Barsh and Schwartz, 2002). In insects, two independent homologs of NPY, neuropeptide F (NPF) and short neuropeptide F (sNPF) (Brown et al., 1999; Hewes and Taghert, 2001), promote feeding behavior (Lee et al., 2004; Wu et al.,

2003) when broadly overexpressed in neurons. Although much is known about the central control of feeding behavior, little is known about starvation modulation of sensory representation in any animal.

For most animals in their natural environment, feeding begins with a search for the appropriate food source, during which the sense of smell plays an indispensable role (Dethier, 1976). Although important inroads have been made in identifying neuropeptides that regulate feeding behavior, little is understood about whether or how these hormones/neuropeptides alter olfaction and how that leads to behavioral changes. In rodents, internal state influences olfactory response in the olfactory cortex (Murakami et al., 2005). However, it is not clear whether these metabolic hormones act directly on the olfactory cortex or whether they play a modulatory role in the olfactory bulb where a variety of different neuromodulators influence neural activity.

Insulin is a global metabolic cue that promotes glucose uptake in both vertebrates and invertebrates (Rulifson et al., 2002). In addition to the regulation of blood glucose, insulin signaling is implicated in the modulation of behaviors relating to feeding, reproduction, and memory (Gerozissis, 2003), and insulin injection into the hypothalamus reduces food intake in rodents (Woods et al., 1998). However, how insulin signaling fine-tunes defined neural circuits to alter behavior is not well understood. Studies of starvation modulation in the *Drosophila* nervous system afford an opportunity to investigate an evolutionarily conserved mechanism for energy homeostasis and establish a causal link between neuropeptide modulation and feeding behavior.

We have investigated whether starvation modulates olfactory processing that mediates food-search behavior. We report that starvation alters olfactory representation of food odor at the first olfactory synapse. The neuropeptide, sNPF, which is expressed in *Drosophila* olfactory receptor neurons (ORNs) (Carlsson et al., 2010; Nassel et al., 2008), mediates this change by facilitating synaptic transmission from select ORNs. Intraglomerular signaling by sNPF is necessary for starvation-dependent enhancement of odor-driven food-search behavior. Furthermore, starvation increases the expression level of the sNPF receptor (sNPFR1) by a reduction of insulin signaling. Thus, neuropeptide signaling causes starvation-dependent presynaptic facilitation of sensory transmission, which optimizes olfactory representation for food finding.

## RESULTS

### Starvation Alters Olfactory Representation and Food-Search Behavior

The antennal lobe is the center for early olfactory processing and is a target for many neuromodulators. Within the antennal lobe, ORNs expressing the same odorant receptor genes (Clyne et al., 1999; Vosshall et al., 1999) converge onto a single glomerulus (Vosshall et al., 2000). ORNs make synapses with many local interneurons and the cognate projection neurons (PNs) (Distler and Boeckh, 1997). Output PNs of the antennal lobe transmit olfactory information from glomeruli to higher brain centers such as the lateral horn and mushroom body (Stocker et al., 1990; Vosshall and Stocker, 2007). Although ORNs are the main drivers of PN output (Olsen et al., 2007; Root et al., 2007), interneurons have been shown to control olfactory sensitivity by presynaptic inhibition (Ignell et al., 2009; Olsen and Wilson, 2008; Root et al., 2008) and lateral excitation (Olsen et al., 2007; Root et al., 2007; Shang et al., 2007). Two neuromodulators, serotonin (Dacks et al., 2009) and tachykinin (Ignell et al., 2009), have been shown to alter antennal lobe activity. If starvation modulates antennal lobe neurons, we should observe a change in odor-evoked activity in PNs.

We performed two-photon imaging to measure PN dendritic calcium responses to odor stimulation in fed and starved flies. Flies bearing *GH146-Gal4* and *UAS-GCaMP* transgenes express the calcium sensor GCaMP in many PNs, allowing the select measurement of calcium response in PN dendrites (Wang et al., 2003). We first imaged responses to apple cider vinegar (Figure 1A), which is highly attractive to *Drosophila* and is a complex odor that resembles a natural food source (Semmelhack and Wang, 2009). Cider vinegar excites five glomeruli at the tested concentrations. Starvation significantly enhances odor response in three glomeruli (DM1, DM4, and DM2) but decreases odor response in two glomeruli (VM2 and VA3; Figures 1B and 1C). It is interesting to note that starvation alters the amplitude of calcium activity without changing the temporal kinetics (Figure 1B). In sharp contrast, our previous study shows that activation of GABA<sub>B</sub> receptors causes presynaptic inhibition and alters the temporal kinetics of PN calcium activity (Root et al., 2008). Therefore, a change in GABA<sub>B</sub> receptor signaling is unlikely to account for the starvation-dependent change in olfactory response. Rather, our results are more consistent with an excitability change in antennal lobe neurons.

The apparent starvation-dependent change of olfactory response in the DM1, DM2, DM4, VM2, and VA3 glomeruli could be due to intra- or interglomerular mechanisms. We therefore investigated starvation modulation of individual glomeruli with reduced lateral activity. To do this, we imaged PN responses to a panel of four different odorants, each of which excites one or two glomeruli at low concentrations (Figures 1D and 1E). The responses in DM1 and DM4 to ethyl acetate were significantly enhanced by starvation, while the response of DM2 but not VM2 to ethyl hexanoate was enhanced by starvation. In contrast, the responses of VA3 and VM2 to 2-phenylethanol and 3-heptanol, respectively, were not modulated by starvation. Therefore, DM1, DM4, and DM2 are more sensitive to odor stimulation in starved animals. However, VA3 and VM2 are not


subject to direct starvation modulation. This result suggests that the apparent suppression of VA3 and VM2 in response to cider vinegar is due to lateral inhibition. We conclude that some antennal lobe neurons are subject to starvation modulation in a glomerular- rather than odor-specific manner, which results in an alteration of the odor map.

Starvation as an internal state affects feeding behavior (Gelperin, 1971), which begins with an olfaction-dependent search for an appropriate food source. Therefore, we expect that the starvation-dependent change in olfactory representation should be matched by an alteration in behavior. We developed a single fly assay that allows the assessment of starvation modulation on odor-driven food-search behavior. We reasoned that latency to find food is a metric of food search. We employed an automatic computer system to monitor the position of individual flies from which we measured the latency required for individuals to reach an odor target. Individual flies were introduced into small arenas each of which contained a food odor, apple cider vinegar, at the center. During the 10 min observation period, starved flies spend most of the time walking near the food source, whereas fed flies wander in the entire arena with a preference for the perimeter (Figure 2A). The latency of food finding is significantly decreased upon starvation (Figure 2B) and is independent of walking speed (Figure S1 available online). Furthermore, surgical removal of the antennae impairs this behavior (Figure S1). Thus, the sense of smell, mediated by the antennae, is required for food-search behavior, and starvation enhances food finding in *Drosophila*.

What is the time course of the starvation-dependent change in olfactory activity and food-search behavior? We first varied starvation time and measured calcium activity of PN dendrites in response to precise electrical stimulation of the olfactory nerve. Nerve stimulation decreases variability in calcium response compared to odor stimulation and allows for finer control of starvation time because the preparation can be done faster than the odor preparation. Imaging calcium activity in PN dendrites of the DM1 glomerulus, we found that calcium activity increased with starvation duration up to 4 hr. Longer starvation duration for 12 hr did not result in more neuronal response (Figures 2C and 2D). We next varied starvation time and examined the latency of food-search behavior. Similar to the starvation-dependent effect on calcium activity, food finding increases up to 4 hr and is not further increased after 12 hr of starvation (Figure 2E). Thus, the change in antennal lobe activity and food-search behavior occurs within 4 hr of starvation.

### sNPF Signaling in ORNs Mediates Starvation Modulation of Food Search

What is the mechanism by which starvation affects odor-guided behavior? The neuropeptide sNPF promotes feeding behavior (Lee et al., 2004) and is expressed in some ORNs (Carlsson et al., 2010; Nassel et al., 2008). We therefore hypothesized that sNPF signaling in ORNs is responsible for the starvation-dependent enhancement of food-search behavior. We expressed RNAi to knockdown expression of sNPF (Figure S2A for knockdown verification) in ORNs of flies bearing the *Or83b-Gal4* and *UAS-sNPF-RNAi* transgenes, and as a control we expressed *sNPF-RNAi* in PNs of flies bearing *GH146-Gal4* and



### Figure 1. Olfactory Representation in Projection Neurons Is Altered by Starvation

(A) Two-photon imaging of PN calcium activity in response to cider vinegar stimulation on two optical planes of the antennal lobe in fed flies. Grayscale images show antennal lobe structure, and pseudocolored images reveal odor-evoked activity at 0.4% SV (saturated vapor pressure).

(B) Representative traces of fluorescence change over time for the five glomeruli excited by cider vinegar at 0.1% SV.


(C) Peak  $\Delta F/F$  across a range of cider vinegar concentrations for each glomerulus.

(D) PN activity of fed flies in response to pure odorants. (E) Peak  $\Delta F/F$  for each glomerulus. (D and E) Odors were applied at the following concentrations (%SV): 1% ethyl acetate 1:10,000 in mineral oil, 0.1% ethyl hexanoate 1:10,000 in mineral oil, 0.5% 2-phenyl ethanol, and 0.1% 3-heptanol.

(C and E)  $n = 5-10$  for each condition; error bars show standard error of the mean (SEM). \* $p < 0.05$ , \*\* $p < 0.01$ , t test. The flies have *GH146-Gal4* and *UAS-GCaMP*. All starvations were 17–24 hr.

*UAS-sNPF-RNAi* transgenes. We measured the latency of food finding in our behavioral assay and found that indeed starved flies lacking sNPF in ORNs exhibit a significantly longer latency in food finding (Figures 3A and 3B). Interestingly, starved sNPF knockdown flies behave similarly to fed flies (Figure S2D), sug-

gesting that low sNPF signaling mimics the fed state in the antennal lobe. The difference in latency between sNPF knockdown flies and control flies cannot be attributed to a change in locomotor activity (Figure S2B). Furthermore, flies with a *P* element disruption of the first intron of the sNPF gene



**Figure 2. Optimum Food-Search Behavior and Peak Olfactory Sensitivity Are Reached within Four Hours of Starvation**

(A) A food-search assay was used to measure the latency of odor-guided food finding. Grayscale image (left) shows an arena with a food odor, 1% cider vinegar, in the center and a single fly (white arrow). The coordinates of single flies are plotted as a function of time in pseudocolor for a representative fed fly and one starved overnight.

(B) The latency of food search as the cumulative percentage of flies that find the odor source over time.

(C and D) Two-photon imaging of PN calcium activity in the DM1 glomerulus in response to electrical stimulation of the olfactory nerve. (C) Representative traces of fluorescence change over time from the DM1 glomerulus in flies with varied starvation durations. (D) Peak  $\Delta F/F$  normalized to the average response without starvation. Stimulation was 1 ms in duration, 10 V in amplitude, and 4 pulses at 100 Hz.  $n = 5-8$  for each starvation condition. Error bars show SEM. \* $p \leq 0.05$ , t test.

(E) Data from behavioral experiments with varied starvation durations shown as the food-finding percentage normalized to that of the fed state. (B and E)  $n = 53-102$  flies for each condition. Error bars show SEM. \* $p \leq 0.05$ , \*\* $p \leq 0.01$ , z test for proportions. See also Figure S1.


(*sNPF<sup>C00448</sup>*) are similarly impaired in food finding (Figure S2E), although these flies may suffer some growth defect (Lee et al., 2008). Thus, sNPF expression in olfactory receptor neurons mediates the starvation-dependent enhancement of food-search behavior.

Although our findings are in accord with previous work indicating that ORNs express the sNPF peptide (Carlsson et al., 2010; Nassel et al., 2008), the population of neurons that express sNPF1 (Feng et al., 2003), the receptor for sNPF, is not known. In salamanders, the NPY receptor localizes to sensory neurons of the olfactory epithelium (Mousley et al., 2006) and is thus poised for a feedback modulation. In the mammalian hypothalamus, NPY neurons project from the arcuate nucleus to the lateral hypothalamus (Barsh and Schwartz, 2002; Cowley et al., 1999) and are poised for a feedforward modulation. Thus, two possible mechanisms may account for the observed modulatory effects of the neuropeptide: (1) if sNPF1 localizes to ORNs, its peptide may modulate starvation-induced behavior through ORN-ORN feedback modulation, or (2) If sNPF1 localizes to PNs, its peptide may modulate starvation-induced behavior through ORN-PN feedforward modulation. To discriminate between these two possibilities, we expressed RNAi to knockdown sNPF1 (Figure S2A for knockdown verification) in either the ORNs or PNs. We found that expression of *sNPF1-RNAi* in ORNs mimics the effect of the neuropeptide knockdown

(Figures 3C and 3D). In contrast, expression of *sNPF1-RNAi* in the PNs has no effect on food-search behavior. The difference in latency between sNPF1 knockdown and control flies cannot be attributed to a change in locomotor activity (Figure S2C). Furthermore, disruption of sNPF1 by expression of a dominant-negative gene (Lee et al., 2008) in ORNs results in a similar decrease in food finding (Figure S2E). Thus, feedback modulation by sNPF1 expressed in ORNs is necessary for starvation-dependent food search.

### Presynaptic Activity in ORNs Is Modulated by sNPF Signaling

Given that knockdown of sNPF and its receptor in ORNs has a profound effect on starvation-dependent food-search behavior, we reasoned that starvation should alter activity in ORN axon terminals. To investigate this, we imaged odor-evoked activity in ORNs in flies that were fed and flies that were starved overnight. Flies bearing the *Or83b-Gal4* and *UAS-GCaMP* transgenes allow the select measurement of calcium activity in ORN axon terminals (Wang et al., 2003). We observed that cider vinegar activates the same five glomeruli when comparing ORNs (Figure 4A) to PNs (Figure 1A). Three glomeruli (DM1, DM4, and DM2) exhibit significant increases in calcium activity upon starvation, whereas the VM2 glomerulus exhibits significant suppression of response at low odor concentration, and



**Figure 3. Starvation-Dependent Food Search Requires sNPF Signaling in ORNs**

The latency of odor-guided food finding was measured in starved flies with 1% cider vinegar.

(A) The coordinates of single flies for representative control flies (left two plots) and those expressing *sNPF-RNAi* (*sNPFi*) in PNs (third from left) or ORNs (right). (B) The latency of food finding.

(C) The coordinates of two representative control flies (left two plots) and those expressing *sNPF1-RNAi* (*sNPFri*) in PNs (third from left) or ORNs (right).

(D) The latency of food finding.  $n = 64$ –103 flies for each condition.

Error bars show SEM. \* $p < 0.05$ , \*\* $p < 0.01$ , z test for proportions comparing the top three curves to the bottom curve in (B) and (D). See also Figure S2.


the VA3 glomerulus is not affected (Figures 4B and 4C). Thus, starvation alters olfactory representation in sensory neurons, which is largely consistent with the changes observed in the antennal lobe output PNs.

We next asked whether sNPF signaling in ORNs causes the starvation-induced changes in olfactory representation. To investigate this, we imaged ORN response to cider vinegar in starved and fed flies with perturbed sNPF signaling. We found that expression of *sNPF-RNAi* in the ORNs eliminates the effect of starvation such that the olfactory representation in starved flies lacking sNPF mirrors that of fed control flies (Figure 4C). The overlapping curves between control fed flies and starved RNAi flies suggest that the effect of RNAi is specific to sNPF signaling rather than a potential nonspecific effect on neuronal properties. Furthermore, there is no difference between starved and fed sNPF knockdown flies, indicating that sNPF mediates the starvation modulation of ORN activity. In addition, expression of RNAi to knockdown sNPF1 in ORNs similarly eliminates the effect of starvation (Figure S3A). We further investigated whether abolishing sNPF signaling presynaptically in ORNs eliminates the starvation-dependent enhancement in postsynaptic PNs. To do this, we used flies bearing the *GH146-LexA*, *LexAop-GCaMP*, *Or83b-Gal4*, and *UAS-sNPF-RNAi* transgenes. Imaging PN calcium activity in the DM1 glomerulus in the absence of presynaptic sNPF, we found that the effect of starvation is abolished such that PN response in starved flies matches that of fed flies (Figures S3B–S3D). The data suggest that the effect of *sNPF-RNAi* is not due to a nonspecific disruption of synaptic transmission from ORNs. Thus, we conclude that

sNPF signaling causes the change in olfactory representation upon starvation.

#### sNPF Signaling Mediates Presynaptic Facilitation

The above results indicate that starvation enhances activity in ORNs by sNPF signaling, suggesting that the neuropeptide could act to facilitate presynaptic activity. To directly test this hypothesis we asked whether exogenous application of sNPF affects presynaptic calcium activity in ORN terminals. In order to eliminate the contribution of any potential modulation at ORN cell bodies, we removed the antennae and delivered precise electrical stimulation to one olfactory nerve while imaging calcium activity in the ipsilateral antennal lobe. We expressed *sNPF-RNAi* in ORNs to eliminate endogenous sNPF, which may occlude the effect of exogenously applied sNPF. Imaging ORN axon terminals, we find that electrical stimulation of the olfactory nerve elicits a calcium transient that is increased upon sNPF application (Figures 4D–4G). Interestingly, this increase occurs only in starved flies but not in fed flies, suggesting that sNPF1 signaling is upregulated upon starvation. We compared the sensitivity to sNPF between the five glomeruli that respond to cider vinegar and found that the DM1, DM2, and DM4 glomeruli exhibit enhanced activity by the neuropeptide, whereas the VM2 and VA3 glomeruli do not (Figure 4G). This result reveals that ORNs terminating in VM2 and VA3 are not modulated by sNPF, which is consistent with the results we obtained with odor stimulation (Figure 1). Therefore, the suppression of calcium activity in VM2 ORNs (Figure 4B) could be a result of lateral presynaptic inhibition (Olsen and Wilson, 2008; Root



**Figure 4. The sNPF Receptor Is Upregulated upon Starvation and Mediates Presynaptic Facilitation in Sensory Neurons**

(A) Two-photon imaging of ORN axon terminal calcium activity in response to cider vinegar stimulation at 0.4% SV in fed flies.

(B) Representative traces of fluorescence change over time for the five glomeruli excited by 0.1% cider vinegar in control flies (top) and those expressing *sNPF-RNAi* in ORNs (sNPFi) (bottom).

(C) Peak  $\Delta F/F$  across a range of cider vinegar concentrations for each glomerulus.  $n = 10\text{--}12$  for each condition; error bars show SEM.  $*p < 0.05$ , t test comparing starved control to fed control. Control flies have *Or83b-Gal4* and *UAS-GCaMP*, and sNPFi flies also have *UAS-sNPF-RNAi* transgenes.

(D–G) ORN axon terminal calcium activity in response to electrical stimulation of the olfactory nerve before and after application of sNPF. (D) Representative traces of fluorescence change over time from the DM1 glomerulus of fed and starved flies in saline and after addition of 10  $\mu\text{M}$  sNPF. (E) Peak  $\Delta F/F$  before and after sNPF. (F) Percent increase in peak  $\Delta F/F$  after exogenous sNPF addition in DM1. (G) Percent increase in peak  $\Delta F/F$  after sNPF addition in starved flies, for the five glomeruli that respond to cider vinegar.


Stimulation was 1 ms in duration, 10 V in amplitude, and 16 pulses at 100 Hz.  $n = 5\text{--}6$ ; error bars show SEM.  $*p < 0.05$ ,  $**p < 0.01$ ,  $***p < 0.001$ , t test. The flies have *Or83b-Gal4* and *UAS-GCaMP* and *UAS-sNPF-RNAi* transgenes. See also Figure S3.

et al., 2008). Furthermore, the suppression of VA3 PN calcium activity (Figure 1B) could be due to lateral feedforward inhibition (Sachse and Galizia, 2002). Thus, the sNPF peptide and its receptor mediate presynaptic facilitation in starved flies at select glomeruli.

#### sNPF Signaling in DM1 Is Necessary and Sufficient for Starvation-Dependent Food-Search Behavior

The ORNs of the DM1, DM2, and DM4 glomeruli have the ability to respond to exogenous sNPF; however the endogenous

source of the neuropeptide is unclear. The peptide could come from receptor neurons of the same glomerulus or alternatively from neighboring glomeruli. We therefore investigated the inter-versus intraglomerular source of sNPF by knocking down sNPF expression in specific ORNs and imaging ORN activity in all glomeruli. Flies bearing the *Or83b-LexA*, *LexAop-GCaMP*, *UAS-sNPF-RNAi*, and *Or-specific-Gal4* transgenes permit the measurement of calcium activity in the axonal termini of many glomeruli, whereas knockdown of sNPF expression is targeted to one specific glomerulus. We found that knockdown of sNPF



**Figure 5. sNPF Signaling in a Single Glomerulus Is Necessary for Starvation-Dependent Food Search**

(A) Two-photon imaging of ORN axon terminals in flies expressing RNAi to knockdown sNPF expression in the ORNs of individual glomeruli. Peak  $\Delta F/F$  normalized to the average response from fed control flies to 0.2% SV cider vinegar.  $n = 5-6$ , \* $p < 0.05$ , t test. All flies have *Or83b-LexA* and *LexAop-GCaMP*, and where indicated, flies also have the *Or-specific-Gal4* and *UAS-sNPF-RNAi*.


(B) The latency of food finding for starved flies expressing RNAi to knockdown sNPF or sNPFR1 in individual glomeruli. RNAi expression in only the DM1 glomerulus significantly decreases food finding.  $n = 80-195$  flies for each condition, \* $p < 0.05$ , z test for proportions comparing control to *sNPFI* and to *sNPFR1*. Error bars show SEM. See also Figure S4.

expression in Or42b ORNs eliminates starvation modulation in only the cognate DM1 glomerulus without any impact on the ORNs of DM2 or DM4 glomeruli (Figure 5A). Similarly, knockdown of sNPF in Or22a and Or59b ORNs abolished starvation modulation in the ORNs of DM2 and DM4 glomeruli, respectively, without any impact on the other glomeruli (Figure 5A). These results suggest that intraglomerular sNPF peptide is necessary whereas interglomerular sNPF is not sufficient for starvation modulation of olfactory sensitivity.

The above results indicate that intraglomerular sNPF signaling selectively increases activity in only three of the five glomeruli activated by cider vinegar. Given that a previous study has found that not all glomeruli contribute equally to odor-guided behavior (Semmelhack and Wang, 2009), we next asked whether sNPF signaling in individual glomeruli is necessary for food-search behavior. We expressed RNAi to knock down the peptide or the receptor in the DM1, DM2, and DM4 ORNs, which are modulated by sNPF. We found that knockdown of the neuropeptide or its receptor in DM1 ORNs results in significantly decreased food finding in starved flies (Figure 5B). This difference cannot be attributed to a difference in locomotor activity (Figure S4A). Strikingly, knockdown of the neuropeptide or its receptor in the DM2 or DM4 ORNs has no effect on the starvation-dependent food-search behavior (Figure 5B). Expression of the RNAi in the VM2 and VA3 ORNs that are not sensitive to sNPF

signaling does not affect food-search behavior (Figure S4B). These results indicate that sNPF signaling in a single ORN channel is necessary for the starvation-dependent food-search behavior.

It has been observed that sNPF is also expressed in the mushroom body (Nassel et al., 2008), which suggests that starvation modulation in the central nervous system could be important for food-search behavior. We therefore evaluated the contribution of the peripheral modulation by performing gain-of-function experiments in fed flies to determine whether peripheral modulation alone is sufficient to induce starvation-like food-search behavior. We first performed imaging experiments to determine whether overexpression of *sNPFR1* increases odor-evoked calcium activity. We imaged calcium activity in Or42b ORNs in control flies bearing the *Or83b-LexA* and *LexAop-GCaMP* transgenes and overexpression flies that also contained the *Or42b-Gal4* and *UAS-sNPFR1* transgenes. Ectopic expression of *sNPFR1* significantly increases  $\Delta F/F$  in the DM1 glomerulus in fed flies (Figure 6A). Furthermore, this enhanced activity is translated into a shorter latency in food-finding behavior in fed flies as ectopic expression of *sNPFR1* leads to increased food finding (Figure 6B). The data suggest that *sNPFR1* overexpression increases activity of Or42b neurons to produce starvation-like behavior. Is a simple increase in sensitivity sufficient to mimic the behavior? To test this, we artificially increased sensitivity of



**Figure 6. Overexpression of sNPFR1 Is Sufficient to Enhance Activity and Food-Search Behavior**

(A) Two-photon imaging of ORN axon terminals in the DM1 glomerulus of fed flies in response to 0.2% SV cider vinegar. Control flies have the *Or83b-LexA* and *LexAop-GCaMP* transgenes, and experimental flies also bear the *Or42b-Gal4* and *UAS-sNPFR1* transgenes.  $n = 5-6$ ,  $*p < 0.05$ ,  $t$  test.

(B) The latency of food finding in fed flies.  $n = 134-168$ ,  $*p < 0.05$ ,  $z$  test for proportions comparing overexpression flies to three controls.

(C) PN dendritic calcium in the DM1 glomerulus of fed flies in response to 0.2% SV cider vinegar. Control flies have *GH146-LexA* and *LexAop-GCaMP*, and experimental flies also have *Or42b-Gal4* and *UAS-sNPFR1* transgenes.  $n = 5-6$ .

(D) The latency of food finding in fed flies.  $n = 66-81$ . Error bars show SEM.

Or42b neurons by ectopically expressing the bacterial sodium channel (NachBac), which has previously been shown to make *Drosophila* neurons hyperexcitable (Nitabach et al., 2006). Indeed expression of NachBac in Or42b neurons produced starvation-like food finding in fed flies (Figure 6B). Thus, modulation of activity in the Or42b ORNs is both necessary for and sufficient to mimic state-dependent food-search behavior. This result also suggests that sNPFR1 is released even in the fed state. Furthermore, the data suggest that modulation of peripheral olfactory activity makes an important contribution to food-search behavior.


In the above experiments, it appears that sNPFR1 is expressed in fed flies but the receptor is only expressed upon starvation. Therefore, overexpression of the peptide in the fed state should not affect neural activity or behavior. To test this hypothesis, we first overexpressed sNPFR1 in ORNs and imaged the postsynaptic PNs. Indeed overexpression of the peptide did not increase activity in the DM1 glomerulus in fed flies (Figure 6C). Next, overexpression of sNPFR1 in Or42b neurons did not alter food-finding behavior in fed flies (Figure 6D). Therefore, an increase in sNPFR1 expression is not sufficient to sensitize ORNs or produce starvation-like food finding in fed flies.

### Insulin Functions as a Satiety Signal to Suppress sNPFR1 Expression

What is the molecular mechanism to increase ORN sensitivity in starved flies to gate appetitive behavior? We first investigated whether this physiological switch involves gene transcription by performing quantitative RT-PCR. We measured the levels of *sNPF* and *sNPFR1* transcripts in isolated antennae of fed and starved flies relative to a control gene, *rp49* (a ribosomal protein). Interestingly, we found that the level of *sNPFR1* mRNA is increased by approximately 4-fold upon starvation, whereas the level of *sNPF* mRNA does not change (Figure 7B). Although we do not detect a change in *sNPF* mRNA, we cannot rule out the possibilities of starvation-dependent changes in neuropeptide translation or release. Nevertheless, ectopic expression of *sNPFR1* expression is sufficient to induce presynaptic facilitation in fed flies (Figure 6A). Therefore, starvation leads to increased expression of *sNPFR1*, which is sufficient to cause presynaptic facilitation even in the absence of any starvation-dependent change in sNPF.

We next asked, what is the metabolic sensor for ORNs to induce expression of *sNPFR1*? It has been well established that the levels of circulating *Drosophila* insulin-like peptide plummet in the starvation state (Geminard et al., 2009), and that the downstream signaling from the insulin receptor (InR) has the capacity to control gene expression (Edgar, 2006). Furthermore, expression of an insulin receptor has been observed in ORNs of *C. elegans* (Chalasan et al., 2010). We therefore asked whether ORNs express the insulin receptor, by assaying immunoreactivity with InR antiserum in flies that express GFP in ORNs. Indeed many of the Or83b neurons (Figure 7A) and all of the Or42b neurons (Figure S5A) have InR immunoreactivity, indicating that the ORNs projecting to the DM1 glomerulus express InR and therefore could be subject to insulin modulation.

Does InR activity alter the expression of sNPFR1 signaling? We reasoned that ectopic expression of a constitutively active InR (InR-CA) in ORNs should mimic the fed state. We first looked at the starvation-dependent expression of *sNPFR1* transcripts and found that starved flies bearing *Or83b-Gal4* and *UAS-InR-CA* do not exhibit an increase in *sNPFR1* transcripts as measured by qRT-PCR (Figure 7B). Similarly, calcium imaging experiments reveal that expression of InR-CA in ORNs eliminates the sensitivity to exogenous sNPF application in the DM1 glomerulus of starved flies (Figure 7C). In starved control flies, bath application of sNPF enhances the axonal calcium transient evoked by electrical stimulation of the olfactory nerve, while sNPF application has no effect on axonal calcium activity in flies expressing InR-CA in ORNs. This experiment was carried out in the same way as those in Figures 4D-4G. These results predict that starvation should not sensitize Or42b ORNs in these flies with the constitutively active InR. Indeed, calcium imaging experiments show that starvation does not increase olfactory response to cider vinegar in DM1 (Figure 7D). Constitutive activation of InR specifically eliminates the starvation-dependent sensitization because the odor response in fed InR-CA flies is not different from fed controls, indicating that the manipulation does not impair these neurons. Measurement of food-search behavior indicates that the constitutively active InR in most



**Figure 7. Insulin Signaling Modulates Expression of *sNPFR1* and Olfactory Sensitivity**

(A) Antennal tissue with immunoreactivity for the InR and GFP expression under the *Or83b* promoter. Tissue was stained with anti-GFP (green) and anti-InR (red) antibodies.

(B) Quantitative RT-PCR analysis of starvation-induced changes in mRNA expression in the antennae of control flies and flies expressing constitutively active InR (InR-CA) in ORNs (left), and that of flies fed PI3K antagonists relative to those fed only sucrose (right).

(C) The change of ORN terminal calcium activity by bath application of sNPFR in starved flies. The antennal nerve was electrically stimulated before and after sNPFR addition the same way as in the experiments in Figures 4D–4G.  $n = 6-9$ .

(D) PN dendritic response to 0.2% SV cider vinegar in the DM1 glomerulus for control flies and those expressing InR-CA in ORNs.  $n = 5-9$ .

(E) The latency of food-search behavior in starved control flies (black and gray) and those expressing InR-CA in ORNs (blue).  $n = 70-90$  flies.

(F) PN dendritic response to 0.2% SV cider vinegar in the DM1 glomerulus for control flies fed sucrose overnight and those fed sucrose plus 25 nM wortmannin or 30  $\mu$ M LY294002.  $n = 5$  each.

(G) The latency of food-search behavior in flies fed wortmannin and LY294002, and control flies fed sucrose only. sNPFRi flies (orange) contain both *Or83b-Gal4* and *UAS-sNPFR1-RNAi*, whereas control flies (black or gray) represent combined data for flies expressing either transgene alone (control groups are not different from each other).  $n = 60-92$  flies.

(H) Model for starvation modulation of olfactory sensitivity. Error bars indicate SEM. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , t test for (B), (C), (D), and (F), and z test for (E) and (G). See also Figure S5.

ORNs (Figure 7E), or selectively in *Or42b* ORNs (Figure S5B), reduces food finding. Therefore, activation of InR in *Or42b* neurons prevents starvation-dependent presynaptic facilitation and food-search behavior.

We next asked whether blockade of InR downstream signals could mimic the effect of starvation in ORNs. Phosphatidylinositol 3-kinase (PI3K) is a crucial downstream molecule for

insulin control of gene transcription and translation to promote cell growth (Leevers et al., 1996; Weinkove et al., 1999). Chronic blockade of insulin signaling can affect growth; therefore we sought to limit the timing of the perturbation. We hypothesized that pharmacological inhibition of PI3K should mimic the starvation state by preventing InR signaling. Two commonly used antitumor drugs, wortmannin and LY294002, have been shown as effective inhibitors of PI3K (Arcaro and Wymann, 1993; Vlahos et al., 1994). Indeed feeding flies overnight with 4% sucrose plus 25 nM wortmannin or 30  $\mu$ M LY294002 sensitizes olfactory response in the DM1 glomerulus to the same level as that of starved flies and significantly greater than that of flies fed only 4% sucrose (Figure 7F). Do these PI3K antagonists alter ORN *sNPFR1* mRNA levels? Indeed qRT-PCR experiments from isolated antennae revealed that feeding flies

with wortmannin or LY294002 causes a significant increase in *sNPFR1* expression relative to flies fed with 4% sucrose (Figure 7B). Thus, either of these PI3K antagonists causes increased expression of *sNPFR1* in ORNs in addition to sensitized olfactory response. Notably, these two PI3K inhibitors appear to increase peptide mRNA level, which is not observed in starved flies; however increased expression of sNPFR does

not change the behavior (Figure 6D). Therefore, we used epistasis experiments to further investigate the link between the drug-induced increase in *sNPFR1* and the drug-induced olfactory sensitization. Expression of *sNPFR1-RNAi* in ORNs eliminates the drug-induced sensitization (Figure 7F), indicating that the sensitization resulting from blocking insulin signaling depends on *sNPFR1* expression in ORNs. Lastly, we asked whether blocking PI3K induces starvation-like behavior in fed flies. Feeding flies either wortmannin or LY294002 leads to significantly increased food finding in comparison to flies expressing *sNPFR1-RNAi* in Or83b neurons and control flies fed only 4% sucrose (Figure 7G). Expression of *sNPFR1-RNAi* specifically in Or42b neurons also eliminates the starvation-like effect of wortmannin (Figure S5C). These results demonstrate that reduced insulin signaling is necessary and sufficient for starvation-dependent upregulation of *sNPFR1* and the induction of presynaptic facilitation, indicating that InR in ORNs is the metabolic sensor to trigger appetitive behavior (Figure 7H).

## DISCUSSION

We report here that a state of starvation modulates olfactory sensitivity at the first synapse in a form of presynaptic facilitation. Starvation increases *sNPFR1* transcription in ORNs, which is both necessary and sufficient for presynaptic facilitation. It has been well established that fluctuation of insulin is a key metabolic cue to maintain energy homeostasis. This study implicates that a low insulin signal via the PI3K pathway increases *sNPFR1* expression. Interestingly, a subset of glomeruli exhibit starvation-dependent presynaptic facilitation that depends on intraglomerular sNPF signaling, whereas selective knockdown of sNPF or *sNPFR1* in only the DM1 glomerulus affects food-search behavior. This finding corroborates our previous work revealing that the DM1 glomerulus is hardwired for innate odor attraction (Semmelhack and Wang, 2009). Thus, an internal state of starvation, with insulin as a global satiety signal acting on sensory neurons through a local sNPF signal, shifts the odor map. Starvation modulation of the odor map increases the saliency of glomerular activity to match the changing physiological needs of an organism.

The Or42b sensory neurons may be considered as a neural substrate for appetitive choices because they integrate internal and external cues to influence an important innate behavior. In this integration, a highly conserved neuropeptide (Hewes and Taghert, 2001) plays an important role in the peripheral olfactory system. A similar presynaptic facilitation mechanism may exist in vertebrates as well. In an aquatic salamander, NPY has been shown to enhance electrical responses of cells in the olfactory epithelium to a food-related odorant in starved animals (Mousley et al., 2006). In addition, NPY immunoreactivity has been observed in the olfactory epithelium of mouse (Hansel et al., 2001) and zebrafish (Mathieu et al., 2002). In the nematode *C. elegans*, elevated activity levels of an NPY-like receptor cause a change in foraging pattern (Macosko et al., 2009). Our study demonstrates that a fluctuating metabolic cue controls *sNPFR1* levels in Or42b neurons, which in turn modulates appetitive behavior. However, it remains to be determined whether other

ORNs mediate attraction behavior and whether they are subject to sNPF-mediated modulation. Given the ubiquitous use of insulin as a metabolic cue, modulation by NPY/sNPF receptors in the early olfactory system could be a conserved mechanism between different animal species.

The internal state of an organism influences its behavior. There is abundant evidence indicating that the global metabolic cue, insulin, works together with local neuropeptides in specific neural circuits to generate state-dependent behavioral responses. In *Drosophila*, the tolerance of a noxious food source is suppressed by insulin signaling and enhanced by NPF signaling such that these two peptides exert their opposing effects on the same neurons that mediate the behavior (Wu et al., 2005). In the mammalian hypothalamus, expression of the orexigenic NPY is suppressed in the satiety state via insulin signaling (Mayer and Belsham, 2009; Schwartz et al., 1992). Results from our study indicate that olfactory response in the periphery is reduced in the satiety state, in which insulin suppresses *sNPFR1* expression to alter neuronal excitability. Insulin's upstream control over *sNPFR1* expression, however, appears to be specific to select neuronal types. Previous work in *Drosophila* has shown that sNPF signaling exerts upstream control of insulin production in the Dilp2 neuroendocrine cells (Lee et al., 2008). In *C. elegans*, the release of an insulin-like peptide in an interneuron is downstream of a neuropeptide involved in promoting behavioral adaptation to food odors (Chalasanani et al., 2010). Thus, different neuronal subtypes may adopt the same neuropeptides for unique and divergent molecular responses. Peptidergic modulation provides a rich repertoire of functional states for the same neural circuit to meet the demand of different internal states.

Central mechanisms to control appetitive behavior, similar to the well-documented modulation of the hypothalamus by NPY, also appear to be important in *Drosophila*. A recent study demonstrates that appetitive memory requires the NPF receptor in the dopaminergic neurons that innervate specific mushroom body lobes (Krashes et al., 2009). This poses the question: what functions are subserved by starvation modulation of multiple neural substrates? It is interesting to note that sensitization of Or42b ORNs is sufficient to enhance food-search behavior in fed flies. Perhaps central modulation by starvation is not necessary for food-search behavior. Modulation in the periphery may serve to gate an animal's sensitivity to specific food odorants, whereas central modulation may serve to enhance an animal's ability to remember the relevant cues in finding a particular food source.

## EXPERIMENTAL PROCEDURES

### Two-Photon Calcium Imaging

GCaMP imaging was performed as previously described (Root et al., 2008; Wang et al., 2003). In odor experiments, a constant airflow of 1 l/min was applied to the antennae via a pipe of 12 mm diameter. Odor onset was controlled by mixing a defined percentage of carrier air with air redirected through odor bottles (presented as %SV, or saturated vapor pressure) as previously described (Root et al., 2008; Semmelhack and Wang, 2009). Nerve stimulation was performed with a glass suction electrode and an S48 stimulator (Grass, Warwick, RI, USA) as previously described (Root et al., 2008; Wang et al., 2003). Starved flies were starved with water for 17–24 hr.

### Behavior Assay

Single female flies were introduced into chambers that were 60 mm in diameter and 6 mm in height. The chamber was illuminated by 660 nm LEDs. Flies were tracked at 2 Hz with custom software written in Labview (V.8.5, National Instruments), and analysis was performed with Igor Pro (V.6, Wavemetrics, Inc) using a custom macro. Latency is defined as the elapsed time before an individual fly spends more than 5 s within a distance of 5 mm from the odor source, which minimizes false positives due to random entry into the odor zone. Apple cider vinegar was diluted 1:100 in 1% low melting temperature agarose, and 5  $\mu$ l were placed in the center of the chamber. Increased cider vinegar concentration leads to increased food finding with a maximum food finding of about 60% at 25% cider vinegar (data not shown). For all of the presented experiments, we used a concentration of cider vinegar that produces moderate food finding such that it could be modulated up or down. In addition, we observed that 17–24 hr starvation and 4–6 hr starvation produced similar results, consistent with the starvation effect measured by calcium imaging (Figure 2E). Therefore, some experiments were carried out with 4–6 hr starvation and others overnight; controls and experimentals were always treated the same.

### Pharmacology

sNPF peptide, AQRSPSLRLRF-NH<sub>2</sub>, 98% purity (Celtek Peptides) was dissolved in saline to a final concentration of 10  $\mu$ M. Wortmannin and LY294002 (LC Laboratories, Woburn, MA, USA) were dissolved in DMSO at stock concentrations of 10 mM and 50 mM, respectively. Flies were fed overnight in vials containing 4% sucrose solution alone, or that plus 25 nM wortmannin or 30  $\mu$ M LY294002.

### Quantitative RT-PCR

RNA was isolated from antennae of 50 female flies for each sample. The RNeasy kit (QIAGEN) was used to isolate RNA and the reverse transcription was performed using the Retroscrip kit (Ambion) with random decamers. This cDNA was subjected to PCR analysis using SYBR green detection on an iCycler thermocycler (Biorad). All values are the average of four replicates, each of which was measured in triplicate and normalized to an rp49 as a control gene.

### Immunostaining

Antennal sections were obtained by mounting live fly heads in OCT and freezing in a dry ice ethanol bath, and 14 mm thick sections were cut on a cryostat. Slides were immediately fixed with ice-cold 4% paraformaldehyde in 0.1 $\times$  PBS for 10 min. Staining was performed using standard techniques with chick-anti-GFP (Ab13970, Abcam, Cambridge, MA, USA) and rabbit-anti-InR (3021, Cell Signaling Technology, Danvers, MA, USA) at 1:1000 and 1:200, respectively.

### Transgenic Flies

See [Extended Experimental Procedures](#) for a list of fly stocks.

### SUPPLEMENTAL INFORMATION

Supplemental Information includes Extended Experimental Procedures and five figures and can be found with this article online at [doi:10.1016/j.cell.2011.02.008](https://doi.org/10.1016/j.cell.2011.02.008).

### ACKNOWLEDGMENTS

We would like to thank R. Axel, M. Gallo, A. Gelperin, S. Kim, and M. Ramaswami for comments on the manuscript. We thank D. Green for assistance with the fly tracking software, E. Troemel for assistance with quantitative RT-PCR, and Jun Hee Lee for helpful discussion of insulin signaling. This work was supported by research grants from the National Institute of Deafness and other Communication Disorders to J.W.W. (R01DC009597, R21DC010458) and fellowship to C.M.R. (1F31DC009511). C.M.R. and J.W.W. designed experiments, analyzed data, and wrote the manuscript. C.M.R., K.I.K., and A.J. carried out experiments.

Received: October 22, 2010

Revised: January 2, 2011

Accepted: February 3, 2011

Published: March 31, 2011

### REFERENCES

- Arcaro, A., and Wymann, M.P. (1993). Wortmannin is a potent phosphatidylinositol 3-kinase inhibitor: the role of phosphatidylinositol 3,4,5-trisphosphate in neutrophil responses. *Biochem. J.* 296, 297–301.
- Barsh, G.S., and Schwartz, M.W. (2002). Genetic approaches to studying energy balance: perception and integration. *Nat. Rev. Genet.* 3, 589–600.
- Berthoud, H.R. (2002). Multiple neural systems controlling food intake and body weight. *Neurosci. Biobehav. Rev.* 26, 393–428.
- Brown, M.R., Crim, J.W., Arata, R.C., Cai, H.N., Chun, C., and Shen, P. (1999). Identification of a *Drosophila* brain-gut peptide related to the neuropeptide Y family. *Peptides* 20, 1035–1042.
- Carlsson, M.A., Diesner, M., Schachtner, J., and Nassel, D.R. (2010). Multiple neuropeptides in the *Drosophila* antennal lobe suggest complex modulatory circuits. *J. Comp. Neurol.* 518, 3359–3380.
- Chalasan, S.H., Kato, S., Albrecht, D.R., Nakagawa, T., Abbott, L.F., and Bargmann, C.I. (2010). Neuropeptide feedback modifies odor-evoked dynamics in *Caenorhabditis elegans* olfactory neurons. *Nat. Neurosci.* 13, 615–621.
- Clyne, P.J., Warr, C.G., Freeman, M.R., Lessing, D., Kim, J., and Carlson, J.R. (1999). A novel family of divergent seven-transmembrane proteins: Candidate odorant receptors in *Drosophila*. *Neuron* 22, 327–338.
- Cowley, M.A., Pronchuk, N., Fan, W., Dinulescu, D.M., Colmers, W.F., and Cone, R.D. (1999). Integration of NPY, AGRP, and melanocortin signals in the hypothalamic paraventricular nucleus: Evidence of a cellular basis for the adipostat. *Neuron* 24, 155–163.
- Dacks, A.M., Green, D.S., Root, C.M., Nighorn, A.J., and Wang, J.W. (2009). Serotonin modulates olfactory processing in the antennal lobe of *Drosophila*. *J. Neurogenet.* 23, 366–377.
- Destexhe, A., and Marder, E. (2004). Plasticity in single neuron and circuit computations. *Nature* 431, 789–795.
- Dethier, V.G. (1976). *The Hungry Fly: A Physiological Study of the Behavior Associated with Feeding* (Cambridge, MA: Harvard University Press).
- Distler, P.G., and Boeckh, J. (1997). Synaptic connections between identified neuron types in the antennal lobe glomeruli of the cockroach, *Periplaneta americana*: II. Local multiglomerular interneurons. *J. Comp. Neurol.* 383, 529–540.
- Edgar, B.A. (2006). How flies get their size: genetics meets physiology. *Nat. Rev. Genet.* 7, 907–916.
- Feng, G., Reale, V., Chatwin, H., Kennedy, K., Venard, R., Ericsson, C., Yu, K., Evans, P.D., and Hall, L.M. (2003). Functional characterization of a neuropeptide F-like receptor from *Drosophila melanogaster*. *Eur. J. Neurosci.* 18, 227–238.
- Gelperin, A. (1971). Regulation of feeding. *Annu. Rev. Entomol.* 16, 365–378.
- Geminard, C., Rulifson, E.J., and Leopold, P. (2009). Remote control of insulin secretion by fat cells in *Drosophila*. *Cell Metab.* 10, 199–207.
- Gerozissis, K. (2003). Brain insulin: regulation, mechanisms of action and functions. *Cell. Mol. Neurobiol.* 23, 1–25.
- Hansel, D.E., Eipper, B.A., and Ronnett, G.V. (2001). Neuropeptide Y functions as a neuroproliferative factor. *Nature* 410, 940–944.
- Hewes, R.S., and Taghert, P.H. (2001). Neuropeptides and neuropeptide receptors in the *Drosophila melanogaster* genome. *Genome Res.* 11, 1126–1142.
- Ignell, R., Root, C.M., Birse, R.T., Wang, J.W., Nassel, D.R., and Winther, A.M. (2009). Presynaptic peptidergic modulation of olfactory receptor neurons in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 106, 13070–13075.

- Krashes, M.J., DasGupta, S., Vreede, A., White, B., Armstrong, J.D., and Waddell, S. (2009). A neural circuit mechanism integrating motivational state with memory expression in *Drosophila*. *Cell* 139, 416–427.
- Lee, K.S., You, K.H., Choo, J.K., Han, Y.M., and Yu, K. (2004). *Drosophila* short neuropeptide F regulates food intake and body size. *J. Biol. Chem.* 279, 50781–50789.
- Lee, K.S., Kwon, O.Y., Lee, J.H., Kwon, K., Min, K.J., Jung, S.A., Kim, A.K., You, K.H., Tatar, M., and Yu, K. (2008). *Drosophila* short neuropeptide F signaling regulates growth by ERK-mediated insulin signalling. *Nat. Cell Biol.* 10, 468–475.
- Leevers, S.J., Weinkove, D., MacDougall, L.K., Hafen, E., and Waterfield, M.D. (1996). The *Drosophila* phosphoinositide 3-kinase Dp110 promotes cell growth. *EMBO J.* 15, 6584–6594.
- Macosko, E.Z., Pokala, N., Feinberg, E.H., Chalasani, S.H., Butcher, R.A., Clardy, J., and Bargmann, C.I. (2009). A hub-and-spoke circuit drives pheromone attraction and social behaviour in *C. elegans*. *Nature* 458, 1171–1175.
- Mathieu, M., Tagliaferro, G., Bruzzone, F., and Vallarino, M. (2002). Neuropeptide tyrosine-like immunoreactive system in the brain, olfactory organ and retina of the zebrafish, *Danio rerio*, during development. *Brain Res. Dev. Brain Res.* 139, 255–265.
- Mayer, C.M., and Belsham, D.D. (2009). Insulin directly regulates NPY and AgRP gene expression via the MAPK MEK/ERK signal transduction pathway in mHypoE-46 hypothalamic neurons. *Mol. Cell. Endocrinol.* 307, 99–108.
- Mousley, A., Polese, G., Marks, N.J., and Eisthen, H.L. (2006). Terminal nerve-derived neuropeptide y modulates physiological responses in the olfactory epithelium of hungry axolotls (*Ambystoma mexicanum*). *J. Neurosci.* 26, 7707–7717.
- Murakami, M., Kashiwadani, H., Kirino, Y., and Mori, K. (2005). State-dependent sensory gating in olfactory cortex. *Neuron* 46, 285–296.
- Nassel, D.R., Enell, L.E., Santos, J.G., Wegener, C., and Johard, H.A. (2008). A large population of diverse neurons in the *Drosophila* central nervous system expresses short neuropeptide F, suggesting multiple distributed peptide functions. *BMC Neurosci.* 9, 90.
- Nitabach, M.N., Wu, Y., Sheeba, V., Lemon, W.C., Strumbos, J., Zelensky, P.K., White, B.H., and Holmes, T.C. (2006). Electrical hyperexcitation of lateral ventral pacemaker neurons desynchronizes downstream circadian oscillators in the fly circadian circuit and induces multiple behavioral periods. *J. Neurosci.* 26, 479–489.
- Olsen, S.R., and Wilson, R.I. (2008). Lateral presynaptic inhibition mediates gain control in an olfactory circuit. *Nature* 452, 956–960.
- Olsen, S.R., Bhandawat, V., and Wilson, R.I. (2007). Excitatory interactions between olfactory processing channels in the *Drosophila* antennal lobe. *Neuron* 54, 89–103.
- Root, C.M., Masuyama, K., Green, D.S., Enell, L.E., Nassel, D.R., Lee, C.H., and Wang, J.W. (2008). A presynaptic gain control mechanism fine-tunes olfactory behavior. *Neuron* 59, 311–321.
- Root, C.M., Semmelhack, J.L., Wong, A.M., Flores, J., and Wang, J.W. (2007). Propagation of olfactory information in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 104, 11826–11831.
- Rulifson, E.J., Kim, S.K., and Nusse, R. (2002). Ablation of insulin-producing neurons in flies: growth and diabetic phenotypes. *Science* 296, 1118–1120.
- Sachse, S., and Galizia, C.G. (2002). Role of inhibition for temporal and spatial odor representation in olfactory output neurons: a calcium imaging study. *J. Neurophysiol.* 87, 1106–1117.
- Schwartz, M.W., Sipols, A.J., Marks, J.L., Sanacora, G., White, J.D., Scheurink, A., Kahn, S.E., Baskin, D.G., Woods, S.C., Figlewicz, D.P., et al. (1992). Inhibition of hypothalamic neuropeptide Y gene expression by insulin. *Endocrinology* 130, 3608–3616.
- Semmelhack, J.L., and Wang, J.W. (2009). Select *Drosophila* glomeruli mediate innate olfactory attraction and aversion. *Nature* 459, 218–223.
- Shang, Y., Claridge-Chang, A., Sjulson, L., Pypaert, M., and Miesenbock, G. (2007). Excitatory local circuits and their implications for olfactory processing in the fly antennal lobe. *Cell* 128, 601–612.
- Stocker, R.F., Lienhard, M.C., Borst, A., and Fischbach, K.F. (1990). Neuronal architecture of the antennal lobe in *Drosophila melanogaster*. *Cell Tissue Res.* 262, 9–34.
- Stocker, R.F., Heimbeck, G., Gendre, N., and de Belle, J.S. (1997). Neuroblast ablation in *Drosophila* P[GAL4] lines reveals origins of olfactory interneurons. *J. Neurobiol.* 32, 443–456.
- Vlahos, C.J., Matter, W.F., Hui, K.Y., and Brown, R.F. (1994). A specific inhibitor of phosphatidylinositol 3-kinase, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). *J. Biol. Chem.* 269, 5241–5248.
- Vosshall, L.B., and Stocker, R.F. (2007). Molecular architecture of smell and taste in *Drosophila*. *Annu. Rev. Neurosci.* 30, 505–533.
- Vosshall, L.B., Amrein, H., Morozov, P.S., Rzhetsky, A., and Axel, R. (1999). A spatial map of olfactory receptor expression in the *Drosophila* antenna. *Cell* 96, 725–736.
- Vosshall, L.B., Wong, A.M., and Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell* 102, 147–159.
- Wang, J.W., Wong, A.M., Flores, J., Vosshall, L.B., and Axel, R. (2003). Two-photon calcium imaging reveals an odor-evoked map of activity in the fly brain. *Cell* 112, 271–282.
- Weinkove, D., Neufeld, T.P., Twardzik, T., Waterfield, M.D., and Leevers, S.J. (1999). Regulation of imaginal disc cell size, cell number and organ size by *Drosophila* class I(A) phosphoinositide 3-kinase and its adaptor. *Curr. Biol.* 9, 1019–1029.
- Woods, S.C., Seeley, R.J., Porte, D., Jr., and Schwartz, M.W. (1998). Signals that regulate food intake and energy homeostasis. *Science* 280, 1378–1383.
- Wu, Q., Wen, T., Lee, G., Park, J.H., Cai, H.N., and Shen, P. (2003). Developmental control of foraging and social behavior by the *Drosophila* neuropeptide Y-like system. *Neuron* 39, 147–161.
- Wu, Q., Zhao, Z., and Shen, P. (2005). Regulation of aversion to noxious food by *Drosophila* neuropeptide Y- and insulin-like systems. *Nat. Neurosci.* 8, 1350–1355.